Maejo International Journal of Science and Technology

ISSN 1905-7873

Available online at www.mijst.mju.ac.th

Review

Chemical constituents and biological activities of *Garcinia* cowa Roxb.

Thunwadee Ritthiwigrom 1,*, Surat Laphookhieo2 and Stephen G. Pyne 3

Received: 2 June 2012 / Accepted: 14 April 2013 / Published: 3 June 2013

Abstract: Garcinia cowa is an abundant source of bioactive phytochemicals. Phytochemical investigations of the plant parts indicated that the fruit, twig and stem are the best source of secondary metabolites, providing flavonoids, phloroglucinols and xanthones respectively. Seventy-eight of these compounds have been identified from the plant and several have interesting pharmacological activities.

Keywords: *Garcinia cowa*, flavonoids, xanthones, phloroglucinols

INTRODUCTION

Many pharmaceutical drug discoveries originated from traditional folk medicine and its associated plant materials and bioactive secondary metabolites. The Genus *Garcinia*, belonging to the Family Clusiaceae which comprises about 300 species, have been widely investigated in terms of their bioactive ingredients. Native to Asia, Africa, South America and Polynesia, the plants are small to medium sized evergreen trees which may grow up to 30 m in height and are widely distributed in the tropical and temperate regions of the world [1]. Twenty-nine species have been observed in Thailand, with 20, 13, 12, 7, 6 and 3 species found in the south, middle, north, east, north-east and west of the country respectively (Figure 1).

Garcinia is a rich source of secondary metabolites, especially triterpenes [1], flavonoids [5], xanthones [6] and phloroglucinols [7]. The latter two groups are well recognised as cheomotaxonomic markers for this genus [8a-e]. Many of the isolated compounds have a wide range

¹Department of Chemistry, Faculty of Science, Chiang Mai University, Chiang Mai, 50200, Thailand

²Natural Products Research Laboratory, School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

³School of Chemistry, University of Wollongong, Wollongong, New South Wales 2522, Australia

^{*} Corresponding author, e-mail: thunwadee.r@cmu.ac.th

of pharmacological activities including anticancer, anti-inflammatory, antibacterial, antiviral, antifungal, anti-HIV, antidepressant and antioxidant [1, 9-13].



Figure 1. Species of *Garcinia* found in different parts of Thailand [2-4]

Garcinia cowa (Figure 2a), commonly known as Cha-muang in Thai, is widely distributed throughout Malaysia, Thailand and Myanmar. The fruits and young leaves are edible with a sour taste. The bark is dark brown with a yellow latex (Figure 2b). The plant has unisex flowers: yellow orange female flowers found at the end of branches and male flowers found along the branches as clusters (Figure 2c). The leaves are glossy, deep green, oblong and up to 6-15 cm in length and 2.5-6.0 cm in width (Figure 2d). The fruits are globose (2.5-6.0 cm in size), green when young and dull

orange or yellow at maturity with 5-8 shallow grooves, at least near the top, and contain 6-8 large 3-angled seeds (Figure 2e) [14].




Figure 2. Parts of *G. cowa*: (a) branch, (b) bark and latex, (c) inflorescences, (d) leaves and (e) ripe and immature fruits (Photos taken by S. Laphookhieo, 2011)

Many parts of *G. cowa* have been used in traditional folk medicine. For example, the bark, latex and root have been used as an antifever agent [15, 16] while the fruit and leaves have been used for indigestion and improvement of blood circulation, and as an expectorant [16]. The chemical composition and biological activities of various parts of *G. cowa* have been investigated. The major compounds found were xanthones and phloroglucinols. However, minor compounds, including depsidones, terpenoids, steroids and flavonoids, were also observed. Currently, 78 compounds have been isolated from the twig [17], stem [18], fruit [19, 20] and latex [15]. This review mainly focuses on the chemical structures and biological activities of the phytochemicals isolated from *G. cowa* and covers the literature up to April 2012.

DISTRIBUTION AND BIOLOGICAL ACTIVITY

The biological activities of the extracts from various parts of *G. cowa* have been investigated, including the hexane and chloroform extracts of the fruit rind and methanol extract of the leaves and twigs [21-23]. The hexane and chloroform extracts from the fruit rind of *G. cowa* were tested against four Gram-positive bacteria (*Bacillus cereus*, *B. coagulans*, *B. subtilis* and *Staphylococcus aureus*) and one Gram-negative bacterium (*Escherichia coli*). Both extracts significantly inhibited bacterial

growth of the Gram-positive bacteria (IC₅₀s 15-30 µg/mL) but not *E. coli* (IC₅₀s 250-500 µg/mL) [21]. The extracts were also found to inhibit the growth of *Aspergillus flavus* ATCC 46283, a common fungal food contaminant which produces aflatoxin B_1 . The degree of inhibition of aflatoxin B_1 production (100% at a concentration of 2000 ppm) was found to be much higher than the inhibition of fungal growth (*ca* 40-60% at the same concentration) [22]. The methanol extracts of the leaves and twigs of *G. cowa* were evaluated for their ability to inhibit low-density lipoprotein peroxidation induced by copper ions. The twig extract had an IC₅₀ value of 20.5 µg/mL and was more potent (higher % inhibition at 1000 µg/mL) than the leaf extract (IC₅₀ not measured). The twig extract was more potent than the leaf extract on platelet aggregation of human whole blood induced by arachidonic acid, adenosine diphosphate and collagen. These activities may be due to the total phenolic content of these extracts, which were 19 and 61 mg of gallic acid equivalent per g of extract for the leaf and twig extracts respectively [23]. The structural types, chemical structures and biological activities of the natural products isolated from different parts of *G. cowa* were summarised in Table 1. A summary of the number of natural product compounds first discovered in each structural class is shown in Figure 3 and from different parts of the plant is shown in Figure 4.

Table 1. Structure, distribution and biological activity of phytochemicals isolated from different parts of *G. cowa*

Category and structure	Name (code), source and reference	Biological activity	
Depsidone			
HO OCH ₃	Cowadepsidone (1); twig [17]*	Cytotoxicity [17] NCI-H187 IC ₅₀ 31.47 μg/mL MFC-7 IC ₅₀ 36.03 μg/mL	
Flavonoid			
HO OH OH OH	R = H Kaempferol (2); branch [24]		
	R = OH Quercetin (3); stem [18]		
ОН	2-(3,5-Dihydroxyphenyl)-2,3-		
	dihy-dro-5,7-dihydroxy-		
HOON	3′,5,5′,7-		
OH O	tetrahydroxyflavanone (4);		
On O	stem [18]		
QH	2-(3,5-Dihydroxy-phenyl)-2,3-		
но	dihy-dro-3,5,7-trihydroxy-		
	(2R,3R)-3,3',5,5',7-		
ОН	pentahydroxy		
OH O	flavanone (5); stem [18]		

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity
Flavonoid		
OH HO, OH OH OH	R = $\frac{7}{2}$ Garccowaside A (6); stem [18]* R = $\frac{7}{2}$ Garccowaside B (7); stem [18]* R = CH ₃ Garccowaside C (8); stem [18]*	
HO OH O	GB-2 (9); branch [24]	
O OH HO OH HO OH	Amentoflavone (10); fruit [19]	
OH OOH OH OH	Morelloflavone (11); fruit [19] twig [16]	Antioxidant activity [16] DPPH assay IC_{50} 10.01 µg/mL Hydroxyl radical scavenging assay IC_{50} 3.11x10 ⁻⁴ µg/mL Superoxide anion scavenging assay IC_{50} 1.50x10 ⁻⁴ µg/mL
HO R ₂	$R_1 = R_2 = H$ Volkensiflavone (12); twig [16]	
OH OH OH OH	$R_1 = \beta$ -glucoside; $R_2 = OH$ Morelloflavone-7'- O -glucoside or fukugiside (13); twig [16]	Antioxidant activity [16] DPPH assay IC_{50} 12.92 $\mu g/mL$ Hydroxyl radical scavenging assay IC_{50} 5.31x10 ⁻⁴ $\mu g/mL$ Superoxide anion scavenging assay IC_{50} 6.39x10 ⁻⁴ $\mu g/mL$

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity
Phloroglucinol		
OH O	Cambogin (14); fruit [25, 26], twig [7], stem [26]	Cytotoxicity [7] against HT-29 HCT-116 CCD-18Co
OH O O O	Guttiferone K (15a) [#] ; fruit [26]*, stem [26]	
HO OH O OH	Guttiferone K (15b)#; twig [7]	Cytotoxicity [7] against HT-29 IC ₅₀ 3.25±0.12 μg/mL HCT-116 CCD-18Co
O OH	Chamuangone (16); leaf [27]*	Antibacterial activity [27] S. pyogenes MIC 7.8 µg/mL S. viridans MIC 15.6 µg/mL H. pylori MIC 15.6 µg/mL B. subtilis, Enterococcus sp. and S. aureus MIC 31.2µg/mL
	Garcicowin A (17); twig [7]*	Cytotoxicity [7] against HT-29 HCT-116 CCD-18Co

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity	
Phloroglucinol			
	$R_1 = $; $R_2 = H$ Garcicowin B (18); twig [7]*	Cytotox against	icity [7] HT-29 HCT-116 CCD-18Co
QH R1		Cytotox	
R ₂ O O O O O O O O O O O O O O O O O O O	$R_1 = CH_3$; $R_2 = OH$ Oblongifolin B (19); twig [7]	against	HT-29 HCT-116 CCD-18Co
	5 <u>~</u> .	Cytotox	
	$R_1 = $; $R_2 = OH$ Oblongifolin C (20); twig [7]	against	HT-29 HCT-116 CCD-18Co
		Cytotox	icity [7]
HO OH O	R = Garcicowin C (21); twig [7]*	against	HT-29 HCT-116 CCD-18Co
	R = Garcicowin D (22); twig [7]*	Cytotox against	
OH O	30-Epicambogin (23); twig [7]	Cytotoxi against IC ₅₀ 3.07	
OH OOH OOH	Oblongifolin A (24); twig [7]		

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity	
Phloroglucinol			
	D =	Cytotox	icity [7]
	$R_1 = \frac{1}{2}$	against	HT-29
	$R_2 =$		HCT-116
HO O O O O	Guttiferone B (25); twig [7]		CCD-18Co
O OH	$R_1 = \frac{e^{\frac{1}{2} \delta_{ij}}}{e^{\frac{1}{2} \delta_{ij}}}$		
	$R_2 = \frac{1}{100}$ Cuttiferone F (26): twig [7]		
	Guttiferone F (26); twig [7]		
QH R ₁	`\$ ² \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	Cytotox	
HO 0 0	$R_1 = R_2 = $	against	HT-29
O OH	Oblongifolin D (27); twig [7]		HCT-116 CCD-18Co
Terpene and Steroid			CCD-18C0
O H H H	Friedelin (28); branch [24]		
HO OH OH	Daucosterol (29); branch [24], fruit [19]		
HO H	β -Sitosterol (30); branch [24], fruit [19]		
HO H	Stigmasterol (31); branch [24]		

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity	
Xanthone			
O OH	R ₁ = OCH ₃ ; R ₂ = H Cowaxanthone A (32); fruit [20] or cowagarcinone C; latex [15]*	Anti-inflammatory activity [16	
O OCH ₃	R ₁ = H; R ₂ = OCH ₃ 1,6-Dihydroxy-3,7-dimethoxy-2- (3-methyl-2-butenyl)xanthone (33); fruit [20]		
	R ₁ = OH; R ₂ = H; R ₃ = R ₄ = OCH ₃ ; R ₅ = prenyl Cowaxanthone B (34); fruit [20]*	Antibacterial activity [20] S. aureus MIC 128 µg/mL MRSA MIC 128 µg/mL Anti-inflammatory activity [16]	
	$R_1 = R_3 = R_4 = OCH_3$; $R_2 = H$; $R_5 = prenyl$ Fuscaxanthone C (35); fruit [20]		
R_4 R_5 R_5 R_5 R_5 R_1 R_2	$R_1 = R_3 = OH$; $R_2 = R_5 = prenyl$; $R_4 = OCH_3$ 7-O-Methylgarcinone E (36); stem [28]*, bark [29], fruit [20]	Antibacterial activity [20] S. aureus MIC 128 µg/mL MRSA MIC 64 µg/mL Antimalarial activity [29] Plasmodium falciparum IC ₅₀ 1.5-3.0 µg/mL	
		Antimalarial activity [29] Plasmodium falciparum IC ₅₀ 1.5-3.0 µg/mL	
	$R_1 = R_3 = OH$; $R_2 = H$; $R_4 = OCH_3$; $R_5 = prenyl$ α -Mangostin (37); fruits [20], bark [29]	Antibacterial activity [20] S.aureus MIC 8 µg/mL MRSA MIC 8 µg/mL Anti-inflammatory activity [16] Antimalarial activity [29] Plasmodium falciparum	
	$R_1 = R_4 = OCH_3$; $R_2 = H$; $R_3 = OH$; $R_5 = PR$ prenyl β -Mangostin (38); fruit [20], twig [17]	IC ₅₀ 1.5-3.0 μg/mL Antibacterial activity [20] S. aureus MIC 128 μg/mL MRSA MIC 64 μg/mL	

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity
Xanthone		
Y	R_1 = prenyl; R_2 = OH Cowaxanthone C (39); fruit [20]*	Antibacterial activity [20] S. aureus MIC 128 µg/mL MRSA MIC 128 µg/mL Anti-inflammatory activity [16]
H_3CO R_2 R_1	$R_1 = H$; $R_2 = OH$ Mangostanin (40); fruit [20]	Antibacterial activity [20] S. aureus MIC 4 µg/mL MRSA MIC 4 µg/mL Anti-inflammatory activity [16]
	R_1 = H; R_2 = OCH ₃ 6-O-Methylmangostanin (41); fruit [20]	
O OH OCH3	Cowaxanthone D (42); fruit [20]*	Anti-inflammatory activity [16]
H ₃ CO OH OH	Cowaxanthone (43); fruit [20], latex [15, 30*], twig [17], bark [29]	Antimalarial activity [29] Plasmodium falciparum $IC_{50} 1.5-3.0 \mu g/mL$ $Cytotoxicity [17]$ NCI-H187 $IC_{50} 3.87 \mu g/mL$ KB $IC_{50} 15.43 \mu g/mL$ MFC-7 $IC_{50} 15.45 \mu g/mL$
	$R_1 = R_3 = \text{prenyl}; R_2 = \text{CHO}$ Cowaxanthone E (44); fruit [20]*	1.0
R_3 O OH R_1 R_2 R_3 O OH R_1	CH ₂ OH R_1 ; $R_2 = H$; $R_3 = geranyl$ Cowanol (45); fruit [20], latex [15, 30*], twig [17], bark [29]	Antibacterial activity [20] S. aureus MIC >128 μ g/mL MRSA MIC >128 μ g/mL Anti-inflammatory activity [16] Antimalarial activity [29] Plasmodium falciparum IC ₅₀ 1.5-3.0 μ g/mL Cytotoxicity [17] NCI-H187 IC ₅₀ 37.26 μ g/mL KB IC ₅₀ 32.34 μ g/mL MFC-7 IC ₅₀ 34.62 μ g/mL

Table 1. (continued)

Category and structure	Name (code), source and reference	Biol	logical activity
Xanthone			
H_3CO H_3C	$R_1 = \text{prenyl}; R_2 = H; R_3 = \text{geranyl}$ Cowanin (46); fruit [20], latex [15, 30*], twig [17], bark [29]	S .aureus MRSA Anti-infl Antin	MIC >128 μg/mL MIC >128 μg/mL MIC >128 μg/mL lammatory activity [16] malarial activity [29] ium falciparum IC ₅₀ 1.5-3.0 μg/mL
R ₄ O OH	R_1 = prenyl; R_2 = OH; R_3 = OCH ₃ ; R_4 = I 1,3,6-Trihydroxy-7-methoxy-2,5- bis(3-methyl-2-butenyl)xanthone (47); latex [15, 30*]	Н	
R_2 OH OH	$R_1 = H$; $R_2 = R_3 = OH$; $R_4 = geranyl$ Norcowanin (48); latex [30]*, twig [17]		Cytotoxicity [17] IC ₅₀ 5.92 μg/mL IC ₅₀ 6.43 μg/mL IC ₅₀ 18.85 μg/mL
R_5 O OH R_1	$R_1 = CH_3$; $R_2 = OH$; $R_3 = prenyl$; $R_4 = OCH_3$; $R_5 = geranyl$ Cowagarcinone A (49); latex [15]*	=	
R_3	$R_1 = CH_3$; $R_2 = R_3 = OCH_3$; $R_4 = R_5 = H$ Cowagarcinone B (50); latex [15]*		
	$R_1 = CH_2OAc;$ $R_2 = OH;$ $R_3 = H;$ $R_4 = OCH_3;$ $R_5 = geranyl$ Cowagarcinone E (51); latex [15]*		
O OH OH	Cowagarcinone D (52); latex [15]*	٠	
ОНОН	Mangostinone (53); latex [15]		
0 OH H ₃ CO	Fuscaxanthone A (54); latex [15]		

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity
Xanthone		
H ₃ CO OH OH OH OH OH OH OH	Cowaxanthone F (55); twig [16]*	
O OH	1,6-Dihydroxyxanthone (56);	
но	twig [16]	
HO O OH H3CO OCH3	3,6-Di- <i>O</i> -methyl-γ-mangostin (57); twig [17]	Cytotoxicity [17] NCI-H187 IC ₅₀ 8.58 μg/mL KB IC ₅₀ 6.64 μg/mL MFC-7 IC ₅₀ 10.59 μg/mL
O OH H ₃ CO OH	Rubraxanthone (58); stem [31, 32]	
0 011	$R_1 = H$; $R_2 = \frac{3}{2}$ 1,5,6-Trihydroxy-3-methoxy-4-(3-hydroxyl-3-methylbutyl)xanthone (59) or nigrolineaxanthone T; stem [18, 33*]	
O OH R_1 HO OCH ₃ R_2	R ₁ = H; R ₂ = prenyl Dulxanthone A (60); stem [18, 33, 34]	Cytotoxicity [34] HepG2
	R_1 = prenyl; R_2 = $\frac{1}{2}$ 4-(1,1-Dimethyl-prop-2-enyl)-1,5,6-trihydroxy-3-methoxy-2-(3-methylbut-2-enyl)xanthen-9-(9 <i>H</i>)-one (61); stem [32]*	
O OH OH R ₂	$R_1 = OCH_3$; $R_2 = prenyl$ 1,5-Dihydroxy-3-methoxy-4-(3-methylbut-2-enyl)-6',6'-dimethyl- 2 <i>H</i> -pyrano(2',3':6,7) xanthone (62); stem [18, 33*]	

Table 1. (continued)


Category and structure	Name (code), source and reference	Biological activity
Xanthone		
O OH OH R ₂	$R_1 = OH; R_2 = H$ 1,3,5-Trihydroxy-6',6'-dimethyl- 2 <i>H</i> -pyrano(2',3':6,7)xanthone (63); stem [18, 33]	
R_4 O OH R_1	$R_1 = R_3 = R_4 = OH; R_2 = H$ 1,3,6,7-Tetrahydroxyxanthone (64) or norathyriol; stem [18, 33] $R_1 = R_2 = OH; R_3 = OCH_3; R_4 = H$ 1,3,5-Trihydroxy-6-methoxy- xanthone (65); stem [33] $R_1 = R_4 = OCH_3; R_2 = R_3 = OH$	
R ₂	1,5,6-Trihydroxy-3,7- dimethoxyxanthone (66); stem [18, 33] $R_1 = R_2 = R_3 = H$; $R_4 = OH$ 1,7-Dihydroxyxanthone (67); stem [33]	
Miscellaneous compound		
OH J J	$(2E,6E,10E)$ -(+)-4 β -Hydroxy-3-	
	methyl-5β-(3,7,11,15- tetramethyl-hexadeca-2,6,10,14- tetraenyl)cyclo-hex-2-en-1-one (68); stem [32]*	
^ ^ ^ ^ ^ ^ ^ ^ ^	O Palmitic acid (69): branch [24]	
	Tetratriacontanoic acid (70); branch [24]	
но он	Oxalic acid (71); fruit [35], fresh leaf [35], bark [35]	
о О ОН ОН	Citric acid (72); fruit [35], fresh leaf [35], bark [35]	
но-Он	4-Hydroxybenzoic acid (73); branch [24]	
H ₃ CO — OH	Isovanillic acid (74); branch [24]	

Table 1. (continued)

Category and structure	Name (code), source and reference	Biological activity
Miscellaneous compound		
НО	<i>p</i> -Coumaric acid (75); fruit [19]	
HO OH	(-)-Hydroxycitric acid lactone (76); fruit [35], fresh leaf [35], bark [35]	
OHC O O CHO	Cirsiumaldehyde (77); fruit [19]	

Note: Biological activity: MRSA (Methicillin resistant *Staphylococcus aureus*), NCI-H187 (Human small cell lung cancer), KB (Oral cavity cancer cell), MCF-7 (Breast cancer cell), HT-29 and HCT-116 (Human colon cancer cell), CCD-18Co (Normal human colon cell), HepG2 (Human hepatocellular liver carcinoma cell)

[#]Guttiferone K (15a, 15b) has been given two different structures in the literature [7, 26].



- □ Number of firstly discovered compounds (new compounds)
- Number of total isolated compounds (new plus known compounds)
- Percentage of total isolated compounds

Figure 3. Classes and numbers of natural products isolated from G. cowa

CLASSESS OF COMPOUNDS ISOLATED FROM G. COWA

Depsidone

Depsidones comprise benzoic acid and phenol skeletons condensed at the *ortho*-positions through ester and ether linkages. This class of natural products is well known in the *Garcinia* species [36, 37]. However, cowadepsidone (1) was the first and only known depsidone from *G. cowa*. It was isolated from the twig extract and showed cytotoxicity against NCI-H187 and MFC-7 cancer cell lines [17].

^{*} Firstly discovered compound (new compound)




Figure 4. Numbers of natural products from different parts of G. cowa

Flavonoids

Twelve flavonoids (compounds **2-13** in Table 1) were isolated from *G. cowa* with garccowasides A **(6)**, B **(7)** and C **(8)** being first reported as new compounds [18]. Of these compounds, only morelloflavone **(11)** and morelloflavone-7"-O-glucoside **(13)** showed strong antioxidant activities [16].

Phloroglucinols

Phloroglucinols are based on a phloroglucinol or 1,3,5-benzenetriol core skeleton or its 1,3,5-cyclohexanetrione (phloroglucin) tautomer. The phloroglucinols found in *G. cowa* have a benzoyl group and geranyl and polyprenyl units as substituent groups. So far, fifteen phloroglucinols (compounds 14-27 in Table 1) have been obtained from the twig including six new compounds: guttiferone K (15a), chamuangone (16), garcicowins A (17), B (18), C (21) and D (22) [7, 26, 27], and nine known phloroglucinols: cambogin (14), guttiferones K (15b), B (25) and F(26), oblongifolins B (19), C (20), A (24) and D (27), and 30-epicambogin (23). Some of them showed selective cytotoxicity against two cancer cell lines (HT-29 and HCT-116) and normal colon cells (CCD-18Co). Guttiferone K (15) and 30-epicambogin (23) exhibited highest cytotoxicity against cancer cell line HT-29 [7]. The name guttiferone K has been given to two different structures in the literature [7, 26] as shown in Table 1. Only one compound, chamuangone (16), was tested for its antibacterial activity and was found to be active against *S. pyogenes* (MIC = 7.8 μ g/mL), *S. viridans* and *H. pylori* (MICs = 15.6 μ g/mL), and *S. aureus*, *B. subtilis* and *Enterococcus* sp. (all of this bacteria shown MICs = 31.2 μ g/mL) [27].

Terpenes and Steroids

Terpenes and steroids represent two large classes of natural products, although they are rare in G. cowa. Only four of these types of compounds (5% of the total compounds isolated) were present in G. cowa, viz. friedelin (28), daucosterol (29), β -sitosterol (30) and stigmasterol (31) [24]. None of these compounds were further studied for their biological activities. However, these compounds which were isolated from other plants had been investigated for their biological

activities. Friedelin (28) from the root bark of Terminalia avicennioides exhibited antibacterial activity against Bacillus Calmette-Guerin (BCG) with an MIC of 4.9 µg/mL [38]. Friedelin (28) and stigmasterol (31) isolated from the leaf of Jatropha tanjorensis were tested against human pathogenic microorganisms, i.e. Gram-positive bacteria: Bacillus cereus, B. subtilis, S. aureus and S. epidermis; Gram-negative bacteria: Aeromonas hydrophila, Escherichia coli, Klebsiella pneumoniae, Pseudomonas aeruginosa, Proteus mirabilis, P. vulgaris, Salmonella paratyphi, S. paratyphi A, Vibrio alcaligenes and V. cholera; and fungi: Aspergillus fumigatus, Candida albicans, Microsporum gypseum and Trichophyton rubrum using the agar-well diffusion and disk diffusion methods [39]. Friedelin (28), at the concentration of 2 µg/mL, showed maximum activity with 37-40, 17-40 and 31-33 mm of clear zone diameter against these three types of microorganisms respectively [39], while stigmasterol (31) at the same concentration exhibited maximum activity with 13-15, 8-17 and 7-8 mm of clear zone diameter respectively [39]. Daucosterol (29) from the roots of Astragalus membranaceus had no growth-inhibitory effect by direct contact but possessed immunomodulatory effect against disseminated candidiasis caused by Candida albicans [40]. β-Sitosterol (30) and stigmasterol (31), isolated from the bark of *Grewia tiliaefolia*, at the same concentration of 1 µg/mL showed antibacterial activity against the Gram-negative bacterium P. aeruginosa (ATCC-20852) with 18 and 20 mm of clear zones respectively and against Klebsiella pneumonia (MTCC-618) with 15 and 15 mm of clear zones respectively as determined by the agar diffusion method [41].

Xanthones

Xanthones, with two aromatic rings linked via carbonyl and ether linkages, are the major components of the *Garcinia* genus [8c-e]. They are commonly found in several parts of *G. cowa*, especially in the stem, fruit and latex. Thirty six xanthones (46% of the total isolated compounds) have been isolated and nineteen of them were first isolated from *G. cowa*. They are cowagarcinone C (32), cowaxanthone (43), cowanol (45), cowanin (46), 1,3,6-trihydroxy-7-methoxy-2,5-bis(3-methyl-2-butenyl)xanthone (47), norcowanin (48), cowagarcinones A (49), B (50), E (51) and D (52) from the latex [15, 30]; cowaxanthones B (34), C (39), D (42) and E (44) from the fruit [20]; 7-*O*-methylgarcinone E (36), 1,5,6-trihydroxy-3-methoxy-4-(3-hydroxyl-3-methylbutyl)xanthone (59), 4-(1,1-dimethyl-prop-2-enyl)-1,5,6-trihydroxy-3-methoxy-2-(3-methylbut-2-enyl)xanthen-9(9*H*)-one (61) and 1,5-dihydroxy-3methoxy-6',6'-dimethyl-2*H*-pyrano(2',3':6,7)-4-(3-methylbut-2-enyl) xanthone (62) from the stem [18, 33]; and cowaxanthone F (55) from the twig [16]. Most of these xanthones showed interesting biological activities.

Antibacterial activity

Eight xanthones from the fruit: cowaxanthones B (34) and C (39), 7-O-methylgarcinone E (36), α -mangostin (37), β -mangostin (38), mangostanin (40), cowanol (45) and cowanin (46) were investigated for their antibacterial activity against *S. aureus* and MRSA. α -Mangostin (37) and mangostanin (40) showed significant activity against these bacteria. α -Mangostin (37) had a MIC value of 8 μ g/mL against both *S. aureus* and MRSA while mangostanin (40) had an MIC value of 4 μ g/mL against both bacteria [20].

Anti-inflammatory activity

Eight xanthones: cowaxanthones A (32), B (34), C (39) and D (42), α -mangostin (37), mangostanin (40), cowanol (45) and cowanin (46) were tested for their anti-inflammatory activity using the ethyl phenylpropiolate induced ear edema assay. All xanthones except cowanol were more active than the standard drug, phenylbutazone [16].

Antimalarial activity

Five xanthones isolated from the stem bark: 7-O-methylgarcinone E (36), α -mangostin (37), cowaxanthone (43), cowanol (45) and cowanin (46) had significant in vitro antimalarial activity against *Plasmodium falciparum* with IC₅₀ values ranging between 1.5-3.0 μ g/mL [29].

Anticancer activity

Six xanthones: cowaxanthone (43), cowanol (45), cowanin (46), norcowanin (48), 3,6-di-O-methyl- γ -mangostin (57) and dulxanthone A (60) isolated from twig were evaluated for their cytotoxicity against NCI-H187, KB, MFC-7 and/or HepG2 cell lines. Cowaxanthone (43), cowanin (46), norcowanin (48) and 3,6-di-O-methyl- γ -mangostin (57) exhibited significant cytotoxicity against the NCI-H187 cell line with IC₅₀ values ranging between 3.87-8.58 µg/mL, and moderately inhibited KB and MCF-7 cancer cell lines with IC₅₀ values ranging between 6.43-15.43 and 10.59-21.38 µg/mL respectively [17]. Dulxanthone A (60) was found to be cytotoxic against the HepG2 cell line [34].

Miscellaneous Compounds

Ten (13% of the total isolated compounds) of the miscellaneous class of compounds have been isolated, including a new discovery: (2E,6E,10E)-(+)-4 β -hydroxy-3-methyl-5 β -(3,7,11,15-tetramethyl-hexadeca-2,6,10,14-tetraenyl)cyclohex-2-en-1-one (68) [32]. None of the isolated compounds from this class were tested for their biological activities.

CONCLUSIONS

 $G.\ cowa$ is an important source of bioactive compounds. Among the parts of this tree, the fruit, twig and stem are the best source of metabolites, thirty of which have been isolated, i.e. one depsidone, one α,β -unsaturated cyclohexenone, three flavonoids, six phloroglucinols and nineteen xanthones. Some of these compounds show interesting pharmacological activities. α -Mangostin (37), cowanol (45) and cowanin (46) are commonly found in all parts of $G.\ cowa$ and they can be used as chemotaxonomic markers of this species. The plant is still under investigation by our research group with the prospect of identifying new bioactive compounds in the near future.

ACKNOWLEDGMENT

The authors thank Chiang Mai University for financial assistance.

REFERENCES

1. A. Kijjoa and L. M. M. Vieira, "Triterpenes from the plants of the Family Clusiaceae (Guttiferae): Chemistry and biological activities", in "Natural Products: Chemistry, Biochemistry and Pharmacology", (Ed. G. Brahmachari), Alpha Science International Ltd., Oxford, **2009**, Ch. 13.

- 2. T. Smitinand, "Thai Plant Names", The Forest Herbarium Royal Forest Department, Bangkok, **2001**, pp.246-248.
- 3. (a) J. F. Maxwell, "Vegetation of Doi Tung, Chiang Rai province, northern Thailand", *Maejo Int. J. Sci. Technol.*, **2007**, *1*, 10-63; (b) J. F. Maxwell, "Addendum to vegetation of Doi Tung, Chiang Rai province, northern Thailand", *Maejo Int. J. Sci. Technol.*, **2008**, *2*, 37-139.
- 4. S. Klaiklay, Y. Sukpondma, V. Rukachaisirikul and S. Phongpaichit, "Friedolanostanes and xanthones from the twigs of *Garcinia hombroniana*", *Phytochem.*, **2013**, *85*, 161-166.
- 5. W. H. Ansari, W. Rahman, D. Barraclough, R. Maynard and F. Scheinmann, "Biflavanoids and a flavanone-chromone from the leaves of *Garcinia dulcis* (Roxb.) Kurz", *J. Chem. Soc., Perkin Trans.1*, **1976**, *13*, 1458-1463.
- 6. V. Rukachaisirikul, T. Ritthiwigrom, A. Pinsa, P. Sawangchote and W. C. Taylor, "Xanthones from the stem bark of *Garcinia nigrolineata*", *Phytochem.*, **2003**, *64*, 1149-1156.
- 7. G. Xu, W. L. T. Kan, Y. Zhou, J.-Z. Song, Q.-B. Han, C.-F. Qiao, C.-H. Cho, J. A. Rudd, G. Lin and H.-X. Xu, "Cytotoxic acylphloroglucinol derivatives from the twigs of *Garcinia cowa*", *J. Nat. Prod.*, **2010**, *73*, 104-108.
- 8. (a) H.-D. Nguyen, B. T.-D. Trinh and L.-H. D. Nguyen, "Guttiferones Q-S, cytotoxic polyisoprenylated benzophenones from the pericarp of *Garcinia cochinchinensis*", *Phytochem. Lett.*, **2011**, *4*, 129-133; (b) S.-C. Chien, C.-F. Chyu, I.-S. Chang, H.-L. Chiu and Y.-H. Kuo, "A novel polyprenylated phloroglucinol, garcinialone, from the roots of *Garcinia multiflora*", *Tetrahedron Lett.*, **2008**, *49*, 5276-5278; (c) Q.-B. Han, N.-Y. Yang, H.-L. Tian, C.-F. Qiao, J.-Z. Song, D. C. Chang, S.-L. Chen, K. Q. Luo and H.-X. Xu, "Xanthones with growth inhibition against HeLa cells from *Garcinia xipshuanbannaensis*", *Phytochem.*, **2008**, *69*, 2187-2192; (d) L.-H. D. Nguyen, H. T. Vo, H. D. Pham, J. D. Connolly and L. J. Harrison, "Xanthones from the bark of *Garcinia merguensis*", *Phytochem.*, **2003**, *63*, 467-470; (e) L.-H. D. Nguyen and L. J. Harrison, "Xanthones and triterpenoids from the bark of *Garcinia vilersiana*", *Phytochem.*, **2000**, *53*, 111-114.
- 9. M. Hemshekhar, K. Sunitha, M. S. Santhosh, S. Devaraja, K. Kemparaju, B. S. Vishwanath, S. R. Niranjana and K. S. Girish, "An overview on genus *Garcinia*: Phytochemical and therapeutical aspects", *Phytochem. Rev.*, **2011**, *10*, 325-351.
- 10. Y. Ren, D. D. Lantvit, E. J. Carcache de Blanco, L. B. S. Kardono, S. Riswan, H. Chai, C. E. Cottrell, N. R. Farnsworth, S. M. Swanson, Y. Ding, X.-C. Li, J. P. J. Marais, D. Ferreira and A. D. Kinghorn, "Proteasome-inhibitory and cytotoxic constituents of *Garcinia lateriflora*: Absolute configuration of caged xanthones", *Tetrahedron*, **2010**, *66*, 5311-5320.
- 11. Y. Sukpondma, V. Rukachaisirikul and S. Phongpaichit, "Xanthone and sesquiterpene derivatives from the fruits of *Garcinia scortechinii*", *J. Nat. Prod.*, **2005**, *68*, 1010-1017.
- 12. H. W. Ryu, M. J. Curtis-Long, S. Jung, Y. M. Jin, J. K. Cho, Y. B. Ryu, W. S. Lee and K. H. Park, "Xanthones with neuraminidase inhibitory activity from the seedcases of *Garcinia mangostana*", *Bioorg. Med. Chem.*, **2010**, *18*, 6258-6264.
- 13. I. O. Pereira, M. J. Marques, A. L. R. Pavan, B. S. Codonho, C. L. Barbieri, L. A. Beijo, A. C. Doriguetto, E. C. D'Martin and M. H. dos Santos, "Leishmanicidal activity of benzophenones and extracts from *Garcinia brasiliensis* Mart. fruits", *Phytomed.*, **2010**, *17*, 339-345.
- 14. T. K. Lim, "Edible Medicinal and Non-Medicinal Plants, Vol. 2: Fruits", Springer Science and Business Media B.V., London, **2012**, pp.29-34.

- 15. W. Mahabusarakam, P. Chairerk and W. C. Taylor, "Xanthones from *Garcinia cowa* Roxb. latex", *Phytochem.*, **2005**, *66*, 1148-1153.
- 16. K. Panthong, N. Hutadilok-Towatana and A. Panthong, "Cowaxanthone F, a new tetraoxygenated xanthone, and other anti-inflammatory and antioxidant compounds from *Garcinia cowa*", *Can. J. Chem.*, **2009**, *87*, 1636-1640.
- 17. S. Cheenpracha, W. Phakhodee, T. Ritthiwigrom, U. Prawat and S. Laphookhieo, "A new depsidone from the twigs of *Garcinia cowa*", *Heterocycles*, **2011**, *83*, 1139-1144.
- 18. J. Shen, Z. Tian and J-s. Yang, "The constituents from the stems of *Garcinia cowa* Roxb. and their cytotoxic activities", *Pharmazie*, **2007**, *62*, 549-551.
- 19. J. Shen and J.-s. Yang, "Chemical constituents from fruit of *Garcinia cowa*", *Zhongguo Yaoxue Zazhi*, **2006**, *41*, 660-661.
- 20. K. Panthong, W. Pongcharoen, S. Phongpaichit and W. C. Taylor, "Tetraoxygenated xanthones from the fruits of *Garcinia cowa*", *Phytochem.*, **2006**, *67*, 999-1004.
- 21. P. S. Negi, G. K. Jayaprakasha and B. S. Jena, "Antibacterial activity of the extracts from the fruit rinds of *Garcinia cowa* and *Garcinia pedunculata* against food borne pathogens and spoilage bacteria", *LWT-Food Sci. Technol.*, **2008**, *41*, 1857-1861.
- 22. G. S. Joseph, G. K. Jayaprakasha, A. T. Selvi, B. S. Jena and K. K. Sakariah, "Antiaflatoxigenic and antioxidant activities of *Garcinia* extracts", *Int. J. Food Microbiol.*, **2005**, *101*, 153-160.
- 23. I. Jantan, F. A. Jumuddin, F. C. Saputri and K. Rahman, "Inhibitory effects of the extracts of *Garcinia* species on human low-density lipoprotein peroxidation and platelet aggregation in relation to their total phenolic contents", *J. Med. Plants Res.*, **2011**, *5*, 2699-2709.
- 24. J. Shen and J. Yang, "Chemical constituents of branch of *Garcinia cowa* Roxb", *Zhongcaoyao*, **2007**, *38*, 993-994.
- 25. G. K. Jayaprakasha, B. S. Jena, L. J. M. Rao and M. C. Varadaraj, "A process for the isolation of cambogin from *Garcinia cowa*", *Indian Pat.*, *No. 242799* (2010).
- 26. J. Shen and J.-S. Yang, "A novel benzophenone from *Garcinia cowa*", *Acta Chim. Sinica*, **2007**, *65*, 1675-1678.
- 27. A. Sakunpak and P. Panichayupakaranant, "Antibacterial activity of Thai edible plants against gastrointestinal pathogenic bacteria and isolation of a new broad spectrum antibacterial polyisoprenylated benzophenone, chamuangone", *Food Chem.*, **2012**, *130*, 826-831.
- 28. K. Lihitwitayawuid, T. Phadungcharoen, C. Mahidol and S. Ruchirawat, "7-O-Methylgarcinone e from *Garcinia cowa*", *Phytochem.*, **1997**, *45*, 1299-1301.
- 29. K. Likhitwitayawuid, T. Phadungcharoen and J. Krungkrai, "Antimalarial xanthones from *Garcinia cowa*", *Planta Med.*, **1998**, *64*, 70-72.
- 30. P. na Pattalung, W. Thongtheeraparp, P. Wiriyachitra and W. C. Taylor, "Xanthones of *Garcinia cowa*", *Planta Med.*, **1994**, *60*, 365-368.
- 31. H.-H. Lee and H.-K. Chan, "1,3,6-Trihydroxy-7-methoxy-8-(3,7-dimethyl-2,6-octadienyl) xanthone from *Garcinia cowa*", *Phytochem.*, **1977**, *16*, 2038-2040.
- 32. F. S. Wahyuni, L. T. Byrne, Dachriyanus, R. Dianita, J. Jubahar, N. H. Lajis and M. V. Sargent, "A new ring-reduced tetraprenyltoluquinone and a prenylated xanthone from *Garcinia cowa*", *Aust. J. Chem.*, **2004**, *57*, 223-226.
- 33. J. Shen and J.-S. Yang, "Two new xanthones from the stems of *Garcinia cowa*", *Chem. Pharm. Bull.*, **2006**, *54*, 126-128.

- 34. Z. Tian, J. Shen, A. P. Moseman, Q. Yang, J. Yang, P. Xiao, E. Wu and I. S. Kohane, "Dulxanthone A induces cell cycle arrest and apoptosis *via* up-regulation of p53 through mitochondrial pathway in HepG2 cells", *Int. J. Cancer*, **2008**, *122*, 31-38.
- 35. B. S. Jena, G. K. Jayaprakasha and K. K. Sakariah, "Organic acids from leaves, fruits, and rinds of *Garcinia cowa*", *J. Agric. Food Chem.*, **2002**, *50*, 3431-3434.
- 36. V. Rukachaisirikul, W. Naklue, S. Phongpaichit, N. H. Towatana and K. Maneenoon, "Phloroglucinols, depsidones and xanthones from the twigs of *Garcinia parvifolia*", *Tetrahedron*, **2006**, *62*, 8578-8585.
- 37. L. D. Ha, P. E. Hansen, F. Duus, H. D. Pham and L.-H. D. Nguyen, "Oliveridepsidones A-D, antioxidant depsidones from *Garcinia oliveri*", *Magn. Reson. Chem.*, **2012**, *50*, 242-245.
- 38. A. Mann, K. Ibrahim, A. O. Oyewale, J. O. Amupitan, M. O. Fatope and J. I. Okogun, "Antimycobacterial friedelane-terpenoid from the root bark of *Terminalia avicennioides*", *Amer. J. Chem.*, **2011**, *1*, 52-55.
- 39. M. B. Viswanathan, J. D. J. Ananthi and P. S. Kumar, "Antimicrobial activity of bioactive compounds and leaf extracts in *Jatropha tanjorensis*", *Fitoterapia*, **2012**, *83*, 1153-1159.
- 40. J.-H. Lee, J. Y. Lee, J. H. Park, H. S. Jung, J. S. Kim, S. S. Kang, Y. S. Kim and Y. Han, "Immunoregulatory activity by daucosterol, a β-sitosterol glycoside, induces protective Th1 immune response against disseminated Candidiasis in mice", *Vaccine*, **2007**, *25*, 3834-3840.
- 41. B. M. K. Ahamed, V. Krishna, H. B. Gowdru, H. Rajanaika, H. M. Kumaraswamy, S. Rajshekarappa, C. J. Dandin and K. M. Mahadevan, "Isolation of bactericidal constituents from the stem bark extract of *Grewia tiliaefolia* Vahl", *Res. J. Med. Plant*, **2007**, *1*, 72-82.

© 2013 by Maejo University, San Sai, Chiang Mai, 50290 Thailand. Reproduction is permitted for noncommercial purposes.